

anti-human CD3 PE-conjugated**Cat-No.: H12125P** **1 ml****Clone:** MEM-57**Specificity:**

The antibody reacts with epsilon and delta chain of CD3 complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of the CD3 subunits CD3gamma, CD3delta, CD3epsilon, and CD3zeta. These CD3 subunits are structurally related members of the immunoglobulin super family encoded by closely linked genes on human chromosome 11. The CD3 components have long cytoplasmic tails that associate with cytoplasmic signal transduction molecules. This association is mediated at least in part by a double tyrosine-based motif present in a single copy in the CD3 subunits. CD3 may play a role in TCR-induced growth arrest, cell survival and proliferation. The CD3 antigen is present on 68-82% of normal peripheral blood lymphocytes, 65-85% of thymocytes and Purkinje cells in the cerebellum. It is never expressed on B or NK cells. Decreased percentages of T lymphocytes may be observed in some autoimmune diseases.

Gene location: human chromosome 11q23

Isotype subclass: Mouse IgG2a**Form:** The purified antibody is conjugated with R-Phycoerythrin (R-PE) under optimum conditions. No reconstitution is necessary.**Expiration date:** The reagent is stable until the expiry date stated on the vial label.**Physical state:** Liquid**Buffer/Additives/Preservative:** PBS containing BSA and 15 mM sodium azide (pH 7.4)**Storage conditions:** Store at 4 °C. Do not freeze. Avoid prolonged exposure to light.**Application:** The reagent is designed for Flow Cytometry analysis of human blood cells.**References:**

Leucocyte Typing III. McMichael A. J. et al (Eds.), Oxford University Press (1987).
Horejsi V. et al., Folia Biol. (Praha) 34, 23 (1988).
Hilgert I. et al., Transplantation 55, 435 (1993).

Warning:

Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research only**. Not for use in human. For in vitro use only.
EuroBioSciences will not be held responsible for patent infringement or other violations that may occur with the use of our products.